

AD-A068 325

TUFTS-NEW ENGLAND MEDICAL CENTER BOSTON MA

F/6 6/5

ISOLATION AND CHARACTERIZATION OF ERYTHROCYTE AND PARASITE MEMB--ETC(U)

APR 78 D F WALLACH

DAMD17-74-C-4118

NL

UNCLASSIFIED

| OF |  
AD  
A068325



END  
DATE  
FILMED  
6-79  
DDC



**LEVEL** *II*

*12*

1.

Isolation and Characterization of Erythrocyte and Parasite  
Membranes from Rhesus Red Cells Infected with P. Knowlesi

III

Annual Summary Report

June 1, 1976 - May 31, 1977

Donald F. H. Wallach

April 1978

Supported by

U. S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND  
Detrick, Frederick, Maryland 21701

Contract No. DAMD 17-74-C-4118

Tufts-New England Medical Center  
171 Harrison Avenue  
Boston, MA 02111



The findings in this report are not to be construed as an  
official Department of the Army position unless so designated  
by other authorized documents

Approved for public release; distribution unlimited.

79 05 03 04 6

ADA068325

DDC FILE COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RESEARCHER'S CATALOG NUMBER	
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED	
Isolation and Characterization of Erythrocyte and Parasite Membranes from Rhesus Red Cells Infected with <u>Plasmodium knowlesi</u> .		Annual <del>Report</del> Summary	repl. no 3
6. AUTHOR(s)		7. PERFORMING ORG. REPORT NUMBER	
Donald F. H. Wallach M. D.		III	
8. PERFORMING ORGANIZATION NAME AND ADDRESS		9. CONTRACT OR GRANT NUMBER(s)	
Tufts-New England Medical Center Boston, Massachusetts 02111		DAMD 17-74-C-4118	
10. CONTROLLING OFFICE NAME AND ADDRESS		11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
US Army Medical Research and Development Command Fort Detrick, Frederick, Maryland 21701		62759A 3A762759A829/00.228	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. REPORT DATE	
(12) 9p.		11 Apr 1978	
		14. NUMBER OF PAGES	
		8	
		15. SECURITY CLASS. (of this report)	
		Unclassified	
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)			
Approved for public release; distribution unlimited.			
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)			
4- A068 308			
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
Pl. knowlesi, erythrocyte-membranes, surface-radiolodination, nitrogen-decompression, differential-centrifugation, dodecyl-sulfate-polyacrylamide-gel-electrophoresis.			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)			
Pure erythrocytes were isolated from normal rhesus monkeys ( <u>Macaca mulatta</u> ) and from animals infected with <u>Plasmodium knowlesi</u> . The external surfaces of both normal and parasitized cells were <sup>125</sup> I-labeled by lactoperoxidase-catalyzed iodination. The erythrocyte membranes were vesiculated in the cold and at physiological ionic strength by nitrogen decompression; 600 psi were used for normal erythrocytes and 280-290 psi for parasitized cells. Host cell membrane and intact, released parasites were isolated by differential centrifugation			

## 20. Abstract ( continued)

and Ficoll-Hypaque density gradient centrifugation; both the yield and degree of purity were good. The fractionations were monitored by (a) microscopy, (b) distribution of erythrocyte membrane-bound  $^{125}\text{I}$ , and (c) analytical dodecyl sulfate polyacrylamide gel electrophoresis. Major differences observed between the membranes of normal and parasitized cells included: (a) presence of an apparently new iodinated glycoprotein component ( Mol wt  $\sim 125,000 - 130,000$ ), (b) apparent monomerization of a glycoprotein normally of Mol wt  $\sim 98,000$ , (c) apparent deletion of an iodinated protein ( Mol wt  $\sim 52,000$ ), and (d) possible selective adsorption to membranes of a parasite protein (Mol wt 68,000).

ACCESSION for	
RTIS	White Section <input checked="" type="checkbox"/>
NDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
SPECIAL	
A	



### Background and Progress Report

To understand the biochemistry and immunology of erythrocytic plasmodial forms, and to develop necessary new chemo- and immunotherapeutic measures for their control requires effective procedures for the separation of parasite and host cell material. Numerous techniques have been applied to the separation of the membranes of parasitized erythrocytes from the enclosed parasites (1-9) but efforts in this direction have been less than satisfactory until very recently. Virtually all of the experimentation in this area has focussed on methods of erythrocyte lysis that might achieve release of intact parasites, namely: osmotic shock (1,2), use of proteases (3), shear by the French press (4), sonication (5) saponin lysis (6,7) and immune lysis (8,9). Indeed, except for the study in the Progress report (10), fractionation criteria in these investigations have been primarily morphological, have not rigorously applied modern technologies for subcellular fractionation (e. g. 11-13), and have essentially neglected the fate of erythrocyte membranes, except to the extent that they constitute morphologically recognizable contaminants of parasite isolates. The latter omission is regrettable in view of the probability that the intraerythrocytic maturation of malaria parasites brings with it not only morphologic (14, 15) biochemical (15) and functional (e. g. 16-18) modifications of the host cell membrane, but immunological alterations as well (e. g. 19).

Abstract

1. Pure erythrocytes are isolated from normal rhesus monkeys (Macaca mulatta) and animals infected with Plasmodium knowlesi.
2. The external surfaces of both cell categories are  $^{125}\text{I}$ -labelled by lactoperoxidase-catalyzed iodination.
3. The erythrocyte membranes are vesiculated in the cold and at physiological ionic strength by nitrogen decompression. 600 psi is used for normal erythrocytes and 280-290 psi for parasitized cells.
4. Host cell membranes and intact, released parasites are isolated in good yield and purity by differential centrifugation and Ficoll-Hypaque density gradient centrifugation.
5. The fractionations are monitored by (a) microscopy (b) distribution of erythrocyte membrane-bound  $^{125}\text{I}$  and (c) analytical dodecyl sulfate polyacrylamide gel electrophoresis.
6. Major differences between the membranes of normal and parasitized cells are observed. These include (a) appearance of an apparently new, iodlatable glycoprotein component (Mol. wt.  $\sim 125,000 - 130,000$ ), (b) apparent monomerization of a glycoprotein normally of Mol. wt.  $\sim 98,000$ , (c) apparent deletion of an iodlatable protein (Mol. wt.  $\sim 52,000$ ), (d) possible selective adsorption to membranes of a parasite protein (Mol. wt. 68,000).

## Progress Report

### Introduction

As part of a program to characterize the host cell membrane modifications that occur during intraerythrocytic maturation of Plasmodium knowlesi, and the different properties of parasite and host cell membranes, we have developed an effective approach for the separation of parasite and host cell components (10). An extension of this approach, as well as detailed characterization of parasite membranes and of parasite-related host cell membrane-protein modifications are the objectives of this application.



### Summary of Proposed Work

1. Continuing initially with the Simian model, the technologies described in item (a) will be modified to give greater cell disruption with lesser parasite damage and better-yield separation of host cell membranes from parasites. For the latter the value of introducing dextran gradients will also be explored.
2. The labelling/monitoring procedures will be extended by techniques for in vitro labelling of host cell membrane sialate and the metabolic labelling of parasite components.
3. The proteins of pure host cell membranes will be further characterized by isoelectric focussing, bidimensional immune electrophoresis, bidimensional focussing - immune electrophoresis and bidimensional DS-PAGE approaches. This involves comparisons with similar analyses on host membranes from normal cells and on parasites, along the lines summarized in (a). The aims here are to (i) define parasite-induced changes in host cell membranes, and (ii) define "candidate" antigens.
4. Techniques currently in use in this laboratory on other systems will be applied to determine conditions required to maintain the permeability characteristics of isolated schizonts.
5. The schizonts isolated after metabolic labelling will be labelled at their external surfaces by lactoperoxidase-catalyzed radioiodination.
6. Conditions will be established to achieve parasite disruption by nitrogen decompression without damage to intracellular organelles.
7. Disrupted parasites will be fractionated into subcellular organelles and soluble fractions, using differential and isopycnic ultracentrifugation and quantitative monitoring using extrinsic and intrinsic markers.
8. Purified membranes will be characterized primarily by the techniques listed in (b3) above.
9. Our approach will be transferred to human in vitro systems when feasible.

1. Christopher, S. R. and Fulton, J. D., *Ann. Trop. Med. Parasitol.* 32 (1938) 43-75.
2. Martin, W. J., Finerty, J. and Rosenthal, A., *Nature New Biol.* 233 (1971) 260-261.
3. Speck, J. F., Moulder, J. W. and Evans, E. A., *J. Biol. Chem.* 164, (1946) 119-144.
4. D'Antonio, L. E., von Doenhoff, A. E., Jr. and Fife, E. H. Jr., *Milit. Med.* 131 (1966) 1152-1156.
5. Kreier, J. P. Pearson, G. L. and Stillwill, D., *Am. J. Trop. Med. Hyg.* 14 (1965) 529-532.
6. Christopher, S. R. and Fulton, J. D., *Ann. Trop. Med. Parasitol.*, 33 (1939) 161-170.
7. Cook R. T., Aikawa M., Rock R. C., Little W. and Spring, H., *Milit. Med.* 134 (1969) 866-883.
8. Langer B. W., Phisphymvidhi P. and Friedlander Y., *Exp. Parasitol.* 20 (1967) 68-76.
9. Shakespeare, P. G. and Trigg, P. I. *Nature (London)*, 241 (1973) 539-540.
10. Wallach, D. F. H. and Conley, M., *J. Mol. Med.* 2, (1977) 119-134.
11. Wallach, D. F. H. and Lin, P. S., *Biochim. Biophys. Acta Reviews on Biomembranes*, 309 (1973) 211-254.
12. Wallach, D.F.H. and Winzler, R., *Evolving Strategies and Tactics in Biomembrane Research*, Chapter 1, Springer Verlag, New York, 1974 pp. 4-56.
13. Wallach, D.F.H. and Schmidt-Ullrich, R., *Methods in Cell Biology*, Vol. 15, (1976) 253-276.
14. Bodammer, J. E. and Bahr, G. F., *Lab. Invest.*, 28 (1973) 708-718.
15. Weidekamm E., Wallach, D.F.H., Hendricks, J. and Lin, P.-S., *Biochim. Biophys. Acta*, 323 (1973) 539-546.
16. Fitch, C. D., Yunis, N. G., Chevli, R. and Gonzalez, Y., *J. Clin. Invest.* 54 (1974) 24-33.
17. Sherman, I. W. and Tanigoshi, L. J., *Protozool.*, 21 (1974) 603-607.
18. Neame, K. D. and Homewood, C. R., *Int. J. Parasitol.*, 5 (1975) 537-540.
19. Wilson, R. J. M., *Parasites in the Immunized Host. Mechanisms of Survival*, Ciba Foundation Symposium, 25 (new series), Excerpta Medica, Amsterdam, 1974, pp. 185-203.

12 Copies

Director (ATTN: SGRD-UWZ-AG)  
Walter Reed Army Institute of  
Research  
Walter Reed Army Medical Center  
Washington, D. C. 20012

4 Copies

HQDA (SGRD-AJ)  
Fort Detrick  
Frederick, MD 21701

12 Copies

Defense Documentation Center  
ATTN: DDC-TCA  
Cameron Station  
Alexandria, Virginia 22314

1 Copy

Dean  
School of Medicine  
Uniformed Services University  
of the Health Sciences  
4301 Jones Bridge Road  
Bethesda, Maryland 20014

1 Copy

Superintendent  
Academy of Health Sciences, US Army  
ATTN: AHS-COM  
Fort Sam Houston, Texas 78234